



Occurrence Details

Occurrence Number: 106C 083

Occurrence Name: Vera Main

Occurrence Type: Hard-rock

Status: Deposit

Date printed: 6/15/2025 11:46:30 AM

General Information

Primary Commodities: lead, silver, zinc

Deposit Type(s): Manto Polymetallic Ag-Pb-Zn, Sediment hosted Mississippi Valley-Type Pb-Zn (MVT)

Location(s): 64°19'3.82" N - -133°45'16.42" W

NTS Mapsheet(s): 106C05

Location Comments: Location based on showings map (AR 093968, pg.10)

Hand Samples Available: Yes

Last Reviewed:

Capsule

Work History

Staked as Vera cl 1-164 (YA37382) in Jul/78 by Prism Joint Venture (Asamera Oil Corporation, Chieftain Development Company Ltd, Prism Resources Ltd, Siebens Oil & Gas Ltd and E & B Exploration Ltd), which carried out geological mapping, geochemical surveying and hand trenching later in the year. In 1979, Dome Petroleum Ltd replaced Siebens and the joint venture carried out geochemical sampling and diamond drilled. In 1980 the joint venture diamond-drilled and carried out 720 m of underground development (drifting and crosscutting). The following year the joint venture carried out underground drilling and in 1982 further prospecting and trenching.

Dome dropped its interest and E & B's was transferred to Imperial Metals Ltd in 1983 and acquired in 1984 by Prism at which time it became project operator. Prism diamond drilled and rock sampled various showings in 1984. In 1988 the company carried out environmental studies for a proposed road into the claims. In 1989 the company purchased a 100% working interest in the Vera claims and the adjoining Val claims to the southeast. In Oct/97, 15966 Yukon Inc a wholly owned subsidiary of Manson Creek Resources Ltd staked Rusty cl 1-131 (YB99989), covering the occurrence. Manson Creek carried out IP geophysical test surveying in 1998 and acquired Landsat satellite imagery.

A regional airborne geophysical survey was conducted over the claim block in 2001 (no report available). Shawn Ryan staked Vera 1-12 claims over the Vera deposit in 2009.

Capsule Geology

The occurrence is located at the southern edge of the Mackenzie Platform, a predominantly shallow water carbonate and clastic sequence that formed on the western margin of the North American craton during Lower Proterozoic through Paleozoic times. The regional geology consists of Upper Proterozoic Rapitan(?) Group mudstones overlain by Upper Proterozoic Profeit Formation dolostones and Upper Proterozoic Nadaleen Formation silty limestone. Over these units are minor clastic and carbonate rocks of the Neoproterozoic to Lower Cambrian Hyland Group. Lower Paleozoic platform carbonates unconformably overlie these units. An arcuate east-west trending, south-dipping normal fault lies north of the occurrence, separating it from Paleoproterozoic Wernecke Supergroup clastic rocks Upper Proterozoic Pinguicula Formation clastics and carbonates to the north.

The Vera deposit is hosted in orange weathering grey argillaceous dolomite of the Profeit Formation (Hay Creek Group). The Main Vera Zone is a vein-fault system within the dolomite with fracture filling of galena, argentiferous tetrahedrite and minor sphalerite, along with dolomite, ankerite, siderite, quartz, limonite, managenese oxides, pyrite, chalcopyrite, scorodite, smithsonite and clay. Mineralization textures range from massive to disseminated sulfides in relatively fresh carbonate gangue to highly oxidized aggregates of limonite, wad, and clay with relict galena and smithsonite. Massive lenses of foliated galena occur in the western part of the Main Zone, and in subsidiary fault zones in the foot and hanging walls. These lenses are narrow and discontinuous but contain substantial quantities of silver.

The Main Zone is variable in total width; it tends to splay towards surface, particularly near the west end, where intervals of mineralized rock occur over true widths of 15 m or more through the zone. At depth, the Main Zone coalesces into two closely spaced well-defined faults. (AR 066208) The system strikes roughly 080 deg and dips roughly 80 deg north. The mineralized vein-fault structure has a strike length of over 655.3 m and a dip length of over 152.4 m (AR 062185).

The best mineralization occurs in two shoots, referred to as West (Vera Main) zone and East zone (Gunsight – MINFILE 106C 114), which are separated by 90 m of lower grade material. The west shoot consists of coarse siderite and dolomite with massive pods of fine-grained galena that are several centimetres to a metre thick along the hanging wall contact, with minor disseminated sphalerite throughout. The east shoot is more brecciated and fractured. The sulfides occur as massive blebs, fine disseminations and tiny veinlets in fractures. Massive and disseminated pyrite occurs and sphalerite is more abundant.

Some of the best surface drill intersections include 1.5 m of 1720 g/t Ag, 59.8% Pb and 0.9% Zn (hole 80-15) and 3.7 m of 1098 g/t Ag, 2.1% Pb and 4.1% Zn (hole 80-17). Drilling in 1984 intersected wider intervals of mineralization, e.g., hole 84-10 – 7.13 m of 205 g/t Ag in the hanging wall and 5.8 m of 1.14% Pb and 72.5 g/t Ag in the footwall (AR 062208).

Underground sampling gave average grades of 696 g/t silver, 6.3% lead and 1.3% zinc across a 3.1 m width and a 131 m length in the West Zone (Vera Main).

Metallurgical and mineralogical testwork in drillcore in 1980 indicate that the silver is present as tetrahedrite-tennantite, both as discrete veins and as intergrowths with galena (AR 062185). Galena and sphalerite were not found to carry silver.

Resource Calculations

Reserves in 1981 were estimated at 628 978 tonnes averaging 303.08 g/t silver, 1.91% lead and 1.75% zinc.

The reserve figures were recalculate in 1982 by the exploration manager at the time giving a 'selective mining reserve' of 352 659 tonnes averaging 606.16 g/t silver, 3.18% lead and 3.47% zinc.

Work History

International Prism Exploration Ltd, p. 18 & 19. Not National Instrument 43-101 compliant. Reported for historical purposes.

1982	MAIN (UNDERGROUND)	Historical Estimate	zinc	3.47 g/t	352,659		No	No	Unknown
Manual calculation by C.M. Lalonde using 1981 underground assay results in conjunction with results of surface and underground drilling.; From Assessment report #062208 by G.W.G. Sivertz for International Prism Exploration Ltd, p. 18 & 19. Not National Instrument 43-101 compliant. Reported for historical purposes.									
1981	MAIN (UNDERGROUND)	Historical Estimate	lead	1.91 %	628,978		No	No	Unknown
Based on calculation by D.H. James, using data from 1979 and 1980 drill holes. Based on 44 drill hole intersections to the -60 m level, with intersections of less than 3 m expanded to 3 m.; From Assessment report #062208 by G.W.G. Sivertz for International Prism Exploration Ltd, p. 17 & 18. Not National Instrument 43-101 compliant. Reported for historical purposes.									
1981	MAIN (UNDERGROUND)	Historical Estimate	silver	303.08 g/t	628,978		No	No	Unknown
Based on calculation by D.H. James, using data from 1979 and 1980 drill holes. Based on 44 drill hole intersections to the -60 m level, with intersections of less than 3 m expanded to 3 m.; From Assessment report #062208 by G.W.G. Sivertz for International Prism Exploration Ltd, p. 17 & 18. Not National Instrument 43-101 compliant. Reported for historical purposes.									
1981	MAIN (UNDERGROUND)	Historical Estimate	zinc	1.75 %	628,978		No	No	Unknown
Based on calculation by D.H. James, using data from 1979 and 1980 drill holes. Based on 44 drill hole intersections to the -60 m level, with intersections of less than 3 m expanded to 3 m.; From Assessment report #062208 by G.W.G. Sivertz for International Prism Exploration Ltd, p. 17 & 18. Not National Instrument 43-101 compliant. Reported for historical purposes.									