



## Occurrence Details

**Occurrence Number:** 105A 017

**Occurrence Name:** Bailey

**Occurrence Type:** Hard-rock

**Status:** Deposit

**Date printed:** 8/6/2025 2:16:26 AM

## General Information

**Primary Commodities:** tungsten, tungsten trioxide

**Secondary Commodities:** copper, molybdenum

**Aliases:** Pat

**Deposit Type(s):** Skarn W

**Location(s):** 60°46'40.8" N - -128°51'3.6" W

**NTS Mapsheet(s):** 105A15

**Hand Samples Available:** No

**Last Reviewed:**

### Capsule

#### Work History

Staked as Pat cl (88383) in Jul/63 and restaked as Tak cl (89115) in May/65 by Cassiar Asbestos Corp L. Restaked as Bailey cl (Y29065) in Mar/69 and as Bailey cl (Y55473) in Jul/71 by Monarch Met ML and Morning Star ML. In Oct/72, Monarch's interest was transferred to Morning Star, which conducted limited hand trenching and sampling in 1971-72. J.C. Turner tied on the Star cl (Y72518) to the southwest in Mar/73 and hand trenched later in the year.

Canada Tungsten Corp optioned the Morning Star claims and explored with mapping, airborne mag and EM surveys and 10 holes (732.4 m) in 1974, about 15 holes in 1975 and 1977, and a legal survey in 1982. Cantung's title was confirmed in a court action between the company and Turner in 1983-84.

O.J. Hall tied on the Pegaseus and Dynamic cl (YA35549) in Sep/78 and conducted hand trenching and mapping in 1979 and 1980, and J. Melnychuk tied on the Peg cl (YA36717) in Jun/79. Paymaster M Inc optioned the Pegaseus and Peg groups and explored with an airborne mag and EM survey and trenching in 1980, and then transferred its interest in 1985 to Cantung, which performed a legal survey later in the year. In July, 1987, the Bailey claims were transferred to T.J.C. Holdings Ltd.

Restaked Apr/92 as part of a group of 90 Logan cl (YB34713) by Placer Dome Inc., which performed reconnaissance prospecting and geochemical sampling.

#### Capsule Geology

Scheelite occurs with pyrrhotite and a trace of chalcopyrite in three separate showings that are exposed over a length of 4 km along the contact between a Cretaceous stock and Devonian-Mississippian limestone. All of the drilling and most other work has been conducted on the B (central) zone, where reserves of 404,600 tonnes grading 1% WO<sub>3</sub> across narrow widths have been outlined. The mineralization is confined to pyroxene-pyrrhotite-scheelite-chalcopyrite skarn near the contact.

The A zone, located 1.6 km north, assayed 0.56% WO<sub>3</sub> across 1.5 m and 0.18% across 10.7 m in a trench 64 m from the intrusive contact. In the C zone, located 2.4 km south, scheelite occurs with molybdenite and pyrrhotite and the best grab sample assayed only 0.06% WO<sub>3</sub>.

In 1992, Placer Dome explored the property for Fort Knox-style gold mineralization, but no significant results were obtained.

### Work History

Date	Work Type	Comment
12/31/1980	Airborne Geophysics	Also EM survey .
12/31/1980	Trenching	
12/31/1975	Drilling	15 holes drilled
12/31/1974	Drilling	Ten holes, 732.4 m.
12/31/1974	Geology	
12/31/1974	Airborne Geophysics	Also EM survey .
12/31/1973	Trenching	
12/31/1971	Geochemistry	
12/31/1971	Trenching	
12/13/1992	Other	
12/13/1985	Other	

### Assessment Reports that overlap occurrence

