

Occurrence Details

Occurrence Number: 116I 084 Occurrence Name: White Fox Occurrence Type: Hard-rock Status: Prospect Date printed: 4/29/2025 12:07:25 AM

General Information

Primary Commodities: vanadium Secondary Commodities: molybdenum, nickel, silver, zinc Deposit Type(s): Sediment hosted Shale-Hosted Ni-Zn-Mo-PGE (Nick) Location(s): 66°54'15.33" N - 136°18'15.41" W NTS Mapsheet(s): 116116 Location Comments: Occurrence location marks best soil sampling result. Hand Samples Available: No Last Reviewed:

Capsule

WORK HISTORY

Staked as Fox cl 1-188 (YC52526) in Feb/2007 by Strategic Metals Ltd. The company immediately optioned the Fox and six other claim groups, collectively called the NiMo (nickel-molybdenum) project to Southampton Ventures Inc, in return for cash, shares and certain work commitments.

During 2007 Southampton Ventures collected 5 rock, 291 soil and 61 silt samples on the Fox claims. The company also drilled 4 diamond drill holes (425.2 m). In Apr/2009 the company changed its name to Quetzal Energy Ltd.

GEOLOGY

The area is located on the west side of the Richardson Trough a north to northwest-trending intracratonic depression formed during Early to Middle Paleozoic time. Deep water shale and argillaceous limestone of the Ordovician to Silurian Road River group are deposited within the trough atop Cambrian and Proterozoic age strata. Younger Paleozoic sediments unconformably cap the Road River Group within the trough and elsewhere in the surrounding broader basin. The entire stratigraphic section is folded by a large-scale anticline that plunges to the north. This anticline is called the Richardson Anticlinorium and its axis approximately coincides with the centre of the trough. To the east, the Richardson Trough is bound by the Trevor Fault and to the West the Deception fault.

The occurrence lies over shallowly (15-20 degrees) west dipping shales which are assigned to the Middle to Upper Devonian Earn Group and Ordovician to Silurian Road River Group. The Earn Group is comprised of sandy shale belonging to the Imperial Formation which conformably overlies siliceous shale of the Canol Formation. Calcareous shale assigned to the Road River Formation unconformably underlies the Canol Formation. The eastern two thirds of the claim block are underlain by Road River shale while the western third is underlain by siliceous shale of the Canol Group. Imperial Formation shale outcrops immediately west of the claim group.

Regionally a thin nickeliferous massive sulphide layer, known as the NiMo horizon is found at the contact between the Canol Formation and Road River Group. This sulphide layer is usually less than 0.5 cm thick and often appears to be lenticular. A similar massive sulphide horizon is reported at the same stratigraphic location at the Nick occurrence (Minfile Occurrence 106D 092) located 245 km to the southwest. The Nick massive sulphide horizon covers an area greater than 80 square kilometres and comprises pyrite, vaesite, melnikovite-type pyrite, sphalerite and wurtzite hosted in a gangue of phosphatic-carbonaceous chert, silica and bitumen. Rock assays from the Nick horizon typically average 3% nickel, 0.20% molybdenum, 0.82% zinc, 0.82% vanadium, 310 ppm platinum and 150 ppb palladium over narrow widths (i.e. < 10 cm). Strategic Metals staked the Fox claims to explore for this type of mineralization.

Southampton Ventures orientated their surface sampling program across the projected surface trace of the Canol Formation – Road River Group contact. Silt samples returned maximum values of 254 ppm nickel, 95.2 ppm molybdenum and 1 230 ppm zinc, while soil samples returned maximum values of 1 255 ppm nickel, 154.5 ppm molybdenum and 2.15% zinc. The highest results were returned from a soil line which crossed a small knoll in the southern part of the claim block (occurrence location).

Diamond drill hole FX07-01 was collared directly into Road River Group rocks while the remaining three holes intersected the Canol Formation - Road River Group contact. All three holes returned only slightly elevated zinc and silver values without any nickel or molybdenum rich massive sulphide layer. The maximum assay values obtained from all core samples was 395 ppm nickel, 2 692 ppm zinc, 66 ppm molybdenum and 2.82 ppm silver. Although no visible mineralization was encountered, it is possible that fine grained tetrahedrite is present in the contact.

Work History

Date	Work Type	Comment
7/1/2020	Geochemistry	
7/1/2020	Geology	
7/1/2020	Geochemistry	
7/1/2020	Other	
7/1/2005	Other	
7/1/2003	Other	
12/31/2007	Geochemistry	
12/31/2007	Drilling	Four holes, 425.2 m.

12/31/2007	Geochemistry	Grid based but lines covered presumed contact.	
12/31/2007	Geochemistry		

Assessment Reports that overlap occurrence

Report Number	Year	Title	Worktypes	Holes Drilled	Meters Drilled
<u>094870</u>	2007	Assessment Report Describing Prospecting, Mapping, Geochemical Sampling and Diamond Drilling at the Fox Property	Diamond - Drilling, Detailed Bedrock Mapping - Geology, Prospecting - Other	4	425.20

Related References

Number	Title	Page(s)	Reference Type	Document Type	
<u>2003-9(D)</u>	Yukon Digital Geology (version 2)		Yukon Geological Survey	Open File (Geological - Bedrock)	
<u>2006-3</u>	Mineral Assessment of the Eagle Plain Study Area, Yukon.		Yukon Geological Survey	Open File (Geological - Bedrock)	
<u>YEG2007 OV</u>	Yukon Exploration and Geology Overview 2007	30, 39, 40.	Yukon Geological Survey	Annual Report	
<u>05-023</u>	Richardson MT Regional Project Report		Yukon Government: Energy, Mines and Resources	YMEP Report	

Drill core at YGS core library

Number	Property	Year Drilled	Core Size	Photos	Data
<u>FX07-01</u>	Fox	2007	BTW	4	1
<u>FX07-02</u>	Fox	2007	BTW	12	3
<u>FX07-03</u>	Fox	2007	BTW	18	4
<u>FX07-04</u>	Fox	2007	BTW	4	2