

Occurrence Details

Occurrence Number: 105F 131
Occurrence Name: Ukon Showing 1
Occurrence Type: Hard-rock

Status: Showing

Date printed: 10/4/2025 1:24:44 AM

General Information

Primary Commodities: niobium, rare earths **Secondary Commodities:** uranium

Aliases: Guano

Deposit Type(s): Skarn

Location(s): 61°29'54.37" N - -132°24'55.71" W

NTS Mapsheet(s): 105F08

Location Comments: Georeferenced from Map 14 in AR 095343 (p. 625).

Hand Samples Available: No

Last Reviewed:

Capsule

Work History

Staked as Guano, etc. cl (YA 00242) in Jul-Sep/76 by Ukon Joint Venture (Chevron and Kerr Addision), which explored with mapping, geochemical sampling and radiometric surveys in 1976. Restaked as PS cl (YB00978) in Aug/87 by Mountain Province Mining Inc which sampled the area, taking contour soil samples downslope of the occurrence and one rock sample in an undisclosed location. In 2003, True North Gems soil sampled in the vicinity of the occurrence. In 2009, True North Gems undertook rock sampling. In 2010, Great Western Minerals carried out a property-wide airborne radiometric & magnetic survey over the property and completed soil lines over the occurrence and to the north of it.

Regional Geology

The occurrence is located on the Cassiar Platform, a curvilinear shelf that formed in the early Paleozoic, roughly parallel to the western margin of the North American craton but separated from it by the Selwyn Basin. Shallow marine miogeoclinal sediments were emplaced on the platform until Late Devonian time. Block faulting and local uplift during the Late Devonian and Mississippian resulted in deposition of carbonaceous shale and chert pebble conglomerate in the Selwyn Basin and across the platform. Local explosive volcanism produced volcaniclastic material and flows of the Pelly Mountains volcanic belt. The belt comprises localized submarine volcanic centres generated in an extensional environment that are separated by basins in-filled with sediments and volcaniclastic rocks. Several cogenetic syenite and trachyte domes and small stocks are the remains of vent areas. Subsequent deformation is a result of Mesozoic thrust faulting related to the Cordilleran orogeny, emplacement of Cretaceous intrusions and Tertiary strike-slip movement along the major northwest-trending Tintina Fault, 30 km to the northeast.

Property Geology

The occurrence is hosted in metasomatized Silurian-Devonian Askin Group carbonates. (It is showing #1 in AR 090269). Chip sampling of the metasomatized carbonate rocks yielded 0.49% total rare earth oxides over 1.0 m. A 30-m-long metre-thick radioactive dyke crosscuts the carbonate and skarn. This resistant east-west trending dyke is fine-grained, green-grey with narrow red/pink bands and quartz veinlets, and numerous carbonate-rich fragments. Despite the high radioactivity, chip sampling of the dyke only returned 0.033% U308. Chip sampling of a dyke 24 m south of the main dyke yielded 1.53% total rare earth oxides, 1.33% ZrO2 and 0.25% Nb2O5 over 0.2 m (AR 095343).

Work History

Date	Work Type	Comment		
12/13/2010	Geochemistry			
12/13/2010	Airborne Geophysics			
12/13/2010	Airborne Geophysics			
12/13/2009	Geochemistry			
12/13/2003	Geochemistry			
12/13/1987	Geochemistry			
12/13/1976	Geochemistry			
12/13/1976	Geology			
12/13/1976	Airborne Geophysics			

Related References

Number	Title	Page(s)	Reference Type	Document Type	
<u>YEG1979 8</u> <u>0-pg55</u>	Rare earth elements in the Guano-Guayes skarn property Pelly Mountains, Yukon Territory		Indian & Northern Affairs Canada/Department of Indian & Northern Development: Exploration & Geological Services Division	Annual Report Paper	
<u>1979Chroni</u> <u>c</u>	Geology of the Guano-Guayes rare earth element bearing skarn property, Pelly Mountains, Yukon Territory		University of British Columbia	MSc Thesis	

Annual Report