



Occurrence Details

Occurrence Number: 105B 150

Occurrence Name: MC Beryl

Occurrence Type: Hard-rock

Status: Showing

Date printed: 12/16/2025 3:20:38 PM

General Information

Secondary Commodities: beryl, gemstones

Deposit Type(s): Gemstone

Location(s): 60°11'33.23" N - -131°37'48.45" W

NTS Mapsheet(s): 105B04

Location Comments: Based on location of beryl sample W10 in AR 094348

Hand Samples Available: No

Last Reviewed:

Capsule

Work History

Staked as Swift 1-8 within a large block of claims in 1978 by the Klinkit Joint Venture (DuPont of Canada Exploration Limited & Duval International Corporation), which explored with geological mapping and contour soil sampling to the north in 1979. The DuPont interest was transferred to CSA Minerals Inc in 1984 and to Goldsearch Inc in 1985. In 1993, H. Kern received a Yukon Mineral Incentives Program (YMIP) grant to explore the area, and prospected and rock sampled that year. I. Elash and Tanana Exploration Inc staked M.C. cl 1-2 (YB93288) in Jul/2001. Elash and Tanana Exploration carried out hand trenching later in the month. The claims were optioned to Strategic Metals Ltd in Aug/2002 which carried out a small hand trenching and prospecting program later in the month. Strategic dropped the option in Nov/2003 and returned the claims to Elash and Tanana Exploration Inc. The claims were optioned to Panarc Resources in 2011, which performed widespread rock sampling of the Seagull Batholith in 2012, although not at the occurrence itself. Panarc Resources staked numerous claim blocks in the area and optioned them to Ucore Rare Metals. In 2014, Ucore flew a large airborne radiometric and magnetic survey over the region, covering almost the entire mapped extent of the Seagull Batholith.

Capsule Geology

This occurrence lies within a belt of Yukon-Tanana Terrane rocks. This belt of rocks is part of an accreted island arc assemblage consisting of bimodal volcanics, coeval plutons and sedimentary rocks, as well as younger Jurassic intrusive rocks and overlap assemblages and Cretaceous intrusions. The occurrence is hosted in mid-Cretaceous Seagull Batholith quartz monzonite.

The occurrence area has predominantly been explored for gem-quality beryl, but exploration in 1993 returned a gold-in-soil value of 288 ppb Au and a rock sample value of 202 ppb Au, 260 m north of the occurrence. However, the author of the report concluded that the anomalies were caused either by small grains of placer gold or by contamination in the laboratory because there were no associated elements in the assay results.

Hand trenching conducted by Elash and Tanana Exploration in 2001 exposed a 20 m section of strongly clay and epidote altered granite containing a partially exposed lens of aggregate intergrown beryl crystals in a saddle between two ridges. Strategic Metals extended the original trench and effectively excavated the beryl lens which measures approximately 80 by 50 by 40 cm in size. An additional lens was exposed 5 m to the south.

The beryl mineralization appears to be associated with multiple acute fault structures that also contain clear quartz flooding and veining. Most of the beryl extracted consists of clear, milky white or pale blue crystal aggregates. Some crystals are up to 7 cm long and 3 cm in diameter, however very few could be segregated without severe damage.

Strategic dug two additional trenches, one on either side of the main trench. The westerly trench encountered beryl float in the upper part of the talus profile but this is believed to be float dispersed down hill from the main showing. Prospecting further along strike to the west uncovered a small amount of beryl float which is also believed to originate from the main showing. No mineralization was found in the easterly trench. South of the main trench prospecting uncovered quartz vein float and float trains which measure up to 25 cm wide and contain 2 to 15 cm vugs and cavities hosting perfectly terminated clear to white quartz crystals and rarer green and purple crystalline fluorite. Strategic Metals felt that since the beryl was hosted within the granite and not on the outer edge where beryllium rich fluids can react with metasedimentary and metavolcanic wall rocks and scavenge sufficient chromium or vanadium, the claims held little potential for emerald mineralization.

Work History

Date	Work Type	Comment
12/13/2014	Airborne Geophysics	
12/13/2014	Airborne Geophysics	
12/13/2002	Trenching	
12/13/2002	Other	
12/13/2001	Trenching	
12/13/1993	Geochemistry	
12/13/1979	Geology	mapping to the north
12/13/1979	Geochemistry	contour soil sampling to the north
12/13/1978	Other	staked

Related References

Number	Title	Page(s)	Reference Type	Document Type
2004-2	Bedrock Geology, Dorsey Lake (NTS 105B/4), southern Yukon (1:50,000 scale)		Yukon Geological Survey	Open File (Geological - Bedrock)
YEG1999-11	Wolf Lake project: Revision mapping of Dorsey Terrane assemblages in the upper Swift River area, southern Yukon and northern B.C.		Indian & Northern Affairs Canada/Department of Indian & Northern Development: Exploration & Geological Services Division	Annual Report Paper
93-043	Report on Prospecting Work in the Dorsey Lake Area		Yukon Government: Energy, Mines and Resources	YMEP Report