



Occurrence Details

Occurrence Number: 115H 072
Occurrence Name: Sack
Occurrence Type: Hard-rock
Status: Showing
Date printed: 6/16/2025 2:33:00 AM

General Information

Primary Commodities: arsenic, gold
Aliases: Kluane, Killer Gold, Ruby Range Project
Deposit Type(s): Unknown
Location(s): N - W
NTS Mapsheet(s): 115H04
Location Comments: Location from map on Strategic Metals Ltd website, May 2022
Hand Samples Available: No
Last Reviewed:

Capsule

The Kluane property is underlain by Late Mesozoic metasedimentary rocks of the Kluane schist and Latest Cretaceous to Eocene rocks of the Ruby Range suite. The Kluane schist consists of a fairly monotonous package of biotite-quartz schist, muscovite-quartz schist and rare lenses of altered and strongly deformed ultramafic rock and marble. Metamorphic grade reaches upper greenschist with local zones of amphibolite. The Kluane schist is intruded by intrusions of the Ruby Range Suite. The Ruby Range suite ranges in age from ca 65 Ma to 52 Ma and consists of biotite granodiorite and hornblende quartz-diorite (Israel et al., 2010).

Three main rock types underlie the Kluane property; 1) biotite-quartz schist of the Kluane schist; 2) migmatitic paragneiss of the Kluane schist; and 3) granodiorite to quartz-diorite of the Ruby Range suite. In the southernmost portion of the claim block, dark grey to black biotite-quartz schist dominates. The schist often forms blocky outcrops that locally weather a brown-orange colour along fractures and joints. Biotite and quartz are the main minerals with the biotite forming the main foliation surfaces. Minor amounts of garnet and staurolite are found as accessory minerals. White to slightly rusty quartz veins of varying widths are ubiquitous and many veins are laterally discontinuous and often have sigmoidal or lozenge shapes. Structurally overlying the schist is a relatively thick package of dark-grey to black and orange weathered migmatitic paragneiss. The paragneiss is characterized by biotite and quartz layers separated by more leucocratic layers comprised of feldspar and quartz. The leucocratic layers appear to be injected melts that may be derived by local partial melting or by melts sourced by intrusions that have migrated along foliation planes.

Marble lenses and rare skarn zones are found within the gneiss on the east-facing slope above Killermun Lake. Exposures of marble are typically white to pale green on both weathered and fresh surfaces, display weak silicification, and range up to 7 m thick and 100 m long. Skarns consist of medium to coarse-grained garnet and diopside. The marbles are often rusty weathering and difficult to distinguish from the surrounding gneiss, unless the rock is broken.

The northern half of the property is comprised of coarse to medium-grained, biotite, granodiorite and hornblende +/- biotite, quartz-diorite of the Ruby Range suite. These rocks generally weather a light grey to white and are locally feldspar porphyritic. Smokey grey quartz grains are common in the granodiorite and less so in the quartz-diorite. The granitic rocks are unfoliated and cross-cut all ductile fabrics found in the Kluane schist except for a strongly foliated border phase that is sporadically observed in outcrop near the boundary with the paragneiss. Foliation in the border phase parallels those found in the Kluane schist and has been attributed to syn-deformation intrusion of the earliest phases of the Ruby Range suite (Israel et al., 2010).

Two sets of narrow unfoliated dykes have been noted on the property in several areas underlain by Kluane schist. The dykes are up to one metre wide and are best distinguished by grain size. One of these dykes, from the central part of the property, has been dated at 55.3 Ma. This overlaps a 55.8 Ma age obtained from the main phase of the Ruby Range found just outside the northern portion of the property.

The Sack Showing lies 1400 m east of the Rikus Zone. It is located on a broad hummock separating two glacial valleys. In 1995, mineralized vein float was discovered along a series of poorly exposed recessive topographic linears. These samples yielded gold values ranging from 3 to 9 g/t (Becker, 1996). In 2014, another two samples were collected from bedrock at the Sack Showing, which returned 6.85 and 2.48 g/t gold (Burrell, 2014). The Sack Showing is arsenopyrite-rich resembles the DalBianco Showing.

Work History

Date	Work Type	Comment
5/1/2020	Geochemistry	
5/1/2020	Geochemistry	
5/1/2020	Trenching	
5/1/2020	Geology	
5/1/2019	Remote Sensing	
5/1/2014	Geochemistry	
5/1/2014	Other	
5/1/2014	Geochemistry	
5/1/2014	Trenching	
5/1/2012	Airborne Geophysics	
5/1/2012	Airborne Geophysics	
5/1/2002	Geochemistry	
5/1/2002	Geochemistry	
5/1/2002	Trenching	
5/1/2002	Other	
5/1/1995	Geochemistry	

5/1/1995	Geology	
5/1/1995	Geochemistry	

Related References				
Number	Title	Page(s)	Reference Type	Document Type
14-076	Assessment Report Describing Prospecting, Hand Trenching and Soil Geochemistry		Yukon Government: Energy, Mines and Resources	YMEP Report